

20. (new) A method for transforming a cell comprising transforming a cell with the isolated nucleic acid fragment of claim 19.

21. (new) A cell comprising the recombinant DNA construct of claim 19.

22. (new) A method for producing a plant comprising transforming a plant cell with the isolated nucleic acid fragment of claim 14 and regenerating a plant from the transformed plant cell.

23. (new) A plant comprising the recombinant DNA construct of claim 19.

24. (new) A seed comprising the recombinant DNA construct of claim 19.

25. (new) An isolated isolated nucleic acid fragment comprising a first nucleotide sequence, wherein the first nucleotide sequence contains at least 30 nucleotides, and wherein the first nucleotide sequence is comprised by another polynucleotide, wherein the other polynucleotide includes:

(a) a second nucleotide sequence, wherein the second nucleotide sequence encodes a polypeptide having methionine synthase activity, wherein the amino acid sequence of the polypeptide and the amino acid sequence of SEQ ID NO:2 or 4 have at least 90% sequence identity based on the Clustal alignment method, or

(b) the complement of the second nucleotide sequence.

26. (new) An isolated polypeptide having methionine synthase activity, wherein the amino acid sequence of the polypeptide and the amino acid sequence of SEQ ID NO: 2 or 4 have at least 90% identity based on the Clustal alignment method.

27. (new) The polypeptide of Claim 26, wherein the amino acid sequence of the polypeptide and the amino acid sequence of SEQ ID NO:2 or 4 have at least 95% identity based on the Clustal alignment method.

28. (new) The polypeptide of Claim 26, wherein the amino acid sequence of the polypeptide comprises the amino acid sequence of SEQ ID NO:2 or 4.

29. (new) A method for isolating a polypeptide encoded by the isolated nucleic acid fragment of claim 14 comprising isolating the polypeptide from a cell containing a recombinant DNA construct comprising the polynucleotide operably linked to at least one regulatory sequence.

30. (new) A nucleic acid fragment comprising

(a) the recombinant DNA construct of claim 19, and

(b) a second recombinant DNA construct comprising a nucleic acid fragment encoding a plant cystathionine γ -synthase or a functionally equivalent subfragment thereof or a complement thereof operably linked to at least one regulatory sequence.

31. (new) A method for increasing methionine content of the seeds of plants comprising:

(a) transforming plant cells with the recombinant DNA construct of claim 19;

(b) growing fertile mature plants from the untransformed plant cells obtained from step (a) under conditions suitable to obtain seeds; and

A1
0001
0002
0003
0004
0005
0006
0007
0008
0009
0010
0011
0012
0013
0014
0015
0016
0017
0018
0019
0020
0021
0022
0023
0024
0025
0026
0027
0028
0029
0030
0031
0032
0033
0034
0035
0036
0037
0038
0039
0040
0041
0042
0043
0044
0045
0046
0047
0048
0049
0050
0051
0052
0053
0054
0055
0056
0057
0058
0059
0060
0061
0062
0063
0064
0065
0066
0067
0068
0069
0070
0071
0072
0073
0074
0075
0076
0077
0078
0079
0080
0081
0082
0083
0084
0085
0086
0087
0088
0089
0090
0091
0092
0093
0094
0095
0096
0097
0098
0099
0100